arXiv:cond-mat/0512605v1 [cond-mat.other] 23 Dec 2005

Mean-field algebraic approach to the dynamics of fermions in a 1D optical lattice

F. P. Massel, V. Penna

Dipartimento di Fisica, Torino Politecnico, Corso Duca degli Abruzzi 24, I-10129 Torino, Italy

E-mail: `francesco.massel@polito.it`

Abstract. We consider a one-dimensional optical lattice of three-dimensional Harmonic Oscillators which are loaded with neutral fermionic atoms trapped into two hyperfine states. By means of a standard variational coherent-state procedure, we derive an effective Hamiltonian for this quantum model and the hamiltonian equations describing its evolution. To this end, we identify the algebra $\mathcal{L}$ of two-fermion operators –describing the relevant microscopic quantum processes of our model– whereby the natural choice for the trial state appears to be a so(2r) coherent state. The coherent-state parameters, playing the role of dynamical variables for the effective Hamiltonian, are shown to identify with the $\mathcal{L}$-operator expectation values thus providing a clear physical interpretation of this algebraic mean-field picture.

PACS numbers: 03.75.Ss, 71.10.Fd, 05.30.Fk, 03.65.Fd

Submitted to: *J. Phys. B: At. Mol. Opt. Phys.*

1. Introduction

Recently, a hierarchy of Hubbard-like Hamiltonians has been proposed to describe the behavior of ultracold fermions in one-dimensional optical lattices (1).

These lattices can be realized with a pair of lasers propagating at a given angle θ ($\theta = \pi$ represents the familiar counterpropagating case), with global confinement ensured by a magnetic trap (see Fig. 1, for a detailed description of this setup see (2)). The pair of lasers give rise to a interference pattern needed to obtain a periodic potential by AC Stark effect. The lattice constant can be adjusted tuning the angle θ according to the relation $d = \lambda/2\sin(\theta/2)$ where λ is the laser wavelength. In addition, it is possible to control both the barrier height of the periodic potential (as a function of the laser intensity) and the interaction between fermions via an external magnetic field (Feshbach resonance, see, e.g. (3)).



Figure 1. Sketch of the experimental setup considered. Two laser beams, propagating at an angle θ, give rise to an AC Stark induced periodic potential. In addition the fermionic cloud is confined by a cigar-shaped magnetic trap with trapping frequencies $\Omega_\perp$ and Ω_x.

These simple considerations allow one to understand how ultracold-atoms physics offers the possibility to explore experimentally a wide range of parameters set that would be unattainable in other contexts, such as the Hubbard model in condensed matter physics.

As a first step towards the description of the cited models, we propose here a mean-field algebraic approach based on coherent-states procedure (4) for a fermionic one-dimensional array of harmonic wells. Although the analytical approach followed here may be regarded as completely general, future numerical analysis will concentrate on a dimer with a six-level structure per well as depicted in Fig. 2). The approach followed here allows one a straightforward reformulation of the the usual mean-field approach for quantum system (based on the 'linearization' of the Hamiltonian and the subsequent solving appropriate self-consistency equations) in terms of a corresponding classical dynamical system. While, for fermions, the interpretation of the aforementioned classical dynamical system as a semiclassical approximation seems not beyond need of justification, it is possible to give a precise physical interpretation to the dynamical variables of the classical problem in terms of expectation values of quantum operators.

In general, it is possible to consider a mean-field approach to a given problem as the constrained minimization of the Hamiltonian $\hat{H}$ over a algebra $\mathcal{L}$. A different choice of $\mathcal{L}$ will lead to different mean-field solutions ((5; 4; 6)). In particular, we will focus on the so(2r) coherent states that, as it will be shown, will lead to the Hartree-Fock-Bogoliubov (7) mean-field approximation, whose effectiveness has been proven for a single spherical harmonic trap in (8).

The paper is organized as follows. In section 2 a brief discussion of the general model considered will be given, along with some possible approximations in different physical situations. As we already mentioned, the fully-analytical control over the physical parameters allows to conceive various Hamiltonians that may have direct experimental relevance. In section 3 so(2r) coherent states and the relevant algebra will be defined. The end of this rather technical section will be devoted to the physical interpretation of the choice of so(2r) as the algebra for the mean-field procedure. In section 4 the classical Hamiltonian $\mathcal{H}_{cl}$ will be deduced and the functional dependence in terms of quantum operators expectation values will be investigated. Finally, in section 4.1 the analysis of the classical dynamical system whose Hamiltonian is $\mathcal{H}_{cl}$ is performed: Lie-Poisson brackets (namely the 'classical' commutators) and, consequently, the evolution equations for the dynamical variables are given.

2. Model Description

In (1), along the lines introduced in (9), a generalized Hubbard Hamiltonian has been introduced to describe the behavior of alkali-metal fermionic atoms in a one-dimensional optical lattice of oblate three dimensional (2+1D) Harmonic Oscillators (pancakes)

$$\hat{\mathsf{H}} = \sum_{\alpha} \lambda_\alpha \hat{n}_\alpha + \sum_{\alpha,\beta} T_{\alpha,\beta} \hat{c}^\dagger_\alpha \hat{c}_\beta + \sum_{\alpha,\beta,\gamma,\delta} U_{\alpha,\beta,\gamma,\delta} \hat{c}^\dagger_\alpha \hat{c}^\dagger_\beta \hat{c}_\delta \hat{c}_\gamma \,. \tag{1}$$

In Eq. (1), α must be considered as a multiple index $\alpha = \{i_\alpha, n_\alpha = 0, J_\alpha, m_\alpha, \sigma_\alpha\}$ whose origin can be traced back to the space(+local)-modes approximation. In this picture n_α, J_α and m_α are the local 2+1 D Harmonic Oscillator quantum numbers, i_α is the site quantum number and σ_α is the spin quantum number. In the following we will confine our analysis to situation where radial modes only are involved in the system dynamics (i.e. we will "freeze" the axial quantum number n_α to zero). The validity of this assumption is guaranteed as long as the radial trapping frequency $\Omega_\perp$ is much smaller than the axial trapping frequency ω_x, i.e. $\Omega_\perp/\omega_x \ll 1$. In this case the tunneling coefficient assumes the following form:

$$T_{\alpha,\beta} = \delta_{J_\alpha,J_\beta} \delta_{m_\alpha,m_\beta} \delta_{\sigma_\alpha,\sigma_\beta} T_{i_\alpha,i_\beta} \tag{2}$$

if we allow nearest-neighbor hopping only

$$T_{i_\alpha,i_\beta} = T\delta_{i_\alpha+1,i_\beta}\ , \tag{3}$$

where T is a known function of the external parameters. Another assumption concerns the two-body interaction term $U_{\alpha,\beta,\gamma,\delta}$, which is treated within the pseudopotential approximation, leading to a delta-like spatial dependence, thus excluding nearest-neighbor two-body interaction terms. If the fermionic nature of the interacting particles is taken into account we have

$$U_{\alpha,\beta,\gamma,\delta} = \delta_{i_\alpha,i_\beta,i_\gamma,i_\delta}\delta_{\sigma_\alpha,\sigma_\gamma}\delta_{\sigma_\beta,\sigma_\delta}\delta_{\sigma_\alpha,-\sigma_\beta}U_{\{J_\alpha,m_\alpha\},\{J_\beta,m_\beta\},\{J_\gamma,m_\gamma\},\{J_\delta,m_\delta\}}\ . \tag{4}$$

Finally, we give the expression for the one-particle energy term which is essentially given by the single-particle energy of the 3D Harmonic Oscillator

$$\lambda_{i_\alpha,n_\alpha,J_\alpha,m_\alpha,\sigma_\alpha} = \left[\hbar\omega_x\left(n_\alpha+\frac{1}{2}\right)+\hbar\Omega_\perp\left(2J_\alpha+1\right)-T_{i_\alpha,i_\alpha}\right]\ , \tag{5}$$

where T_{i_α,i_α} represents a "hopping correction" to the single particle energy term.

For the case considered in Fig. 2 the selection rules imposed on the two-body interaction term select three possible values on $U_{\alpha,\beta,\gamma,\delta}$ that can be classified as: i) lowest-level/ lowest-level interaction terms, ii) lowest-level/highest-level interaction terms, iii) highest-level/highest-level interaction terms.


$\{\imath_\alpha, n_\alpha, J_\alpha, m_\alpha, \sigma_\alpha\} = \{0,0,1/2,1/2,\pm1/2\}$
$\{\imath_\alpha, n_\alpha, J_\alpha, m_\alpha, \sigma_\alpha\} = \{0,0,1/2,-1/2,\pm1/2\}$
$\{\imath_\alpha, n_\alpha, J_\alpha, m_\alpha, \sigma_\alpha\} = \{0,0,0,0,\pm1/2\}$

Figure 2. Schematic view of the single-particle energy levels. Up and down arrows represent possible spin values. We have explicitly indicated single-particle quantum numbers for the left-hand-side well. Analogously, it is possible to define quantum numbers for the right-hand-side well.

3. Coherent States

Since in our mean-field analysis we would like to keep trace of the (possible) atom pairing, the most appropriate coherent-states algebra seems to be, according to (4), the algebra spanned by the $r(2r-1)$ operators $\{\hat{c}^\dagger_\alpha \hat{c}_\beta (1 \le i, j \le r), \hat{c}_\alpha \hat{c}_\beta, \hat{c}^\dagger_\alpha \hat{c}^\dagger_\beta\}$, i.e. so(2r). Its commutation relations can be written as

$$\begin{aligned}
\left[Y^1_{\alpha\beta}, Y^1_{\gamma\delta}\right] &= \left[Y^2_{\alpha\beta}, Y^2_{\gamma\delta}\right] = 0 \\
\left[Y^1_{\alpha\beta}, Y^2_{\gamma\delta}\right] &= Y^3_{\gamma\beta}\delta_{\alpha\delta} + Y^3_{\delta\alpha}\delta_{\beta\gamma} - Y^3_{\gamma\alpha}\delta_{\beta\delta} - Y^3_{\delta\beta}\delta_{\alpha\gamma} \\
\left[Y^1_{\alpha\beta}, Y^3_{\gamma\delta}\right] &= Y^1_{\alpha\delta}\delta_{\beta\gamma} - Y^1_{\beta\delta}\delta_{\alpha\gamma} \\
\left[Y^2_{\alpha\beta}, Y^3_{\gamma\delta}\right] &= Y^2_{\beta\gamma}\delta_{\alpha\delta} - Y^2_{\alpha\gamma}\delta_{\beta\delta} \,,
\end{aligned} \tag{6}$$

having defined

$$Y^1_{\alpha\beta} = \hat{c}_\alpha \hat{c}_\beta \,, \tag{7}$$

$$Y^2_{\alpha\beta} = \hat{c}^\dagger_\beta \hat{c}^\dagger_\alpha \,, \tag{8}$$

$$Y^3_{\alpha\beta} = \hat{c}^\dagger_\alpha \hat{c}_\beta \,. \tag{9}$$

With the above definitions the coherent states can be expressed as

$$|\phi\rangle = \exp\left[- \sum_{1 \le \alpha \ne \beta \le r} \left(\eta_{\alpha,\beta} \hat{c}^\dagger_\alpha \hat{c}^\dagger_\beta - H.c. \right) \right] |0> \ . \tag{10}$$

To evaluate the expectation value of the Hamiltonian $\hat{H}$ defined by equation (1) over the coherent state of the form (10), it is necessary to evaluate the action of the operator

$$\hat{\Omega} = \exp\left[- \sum_{1 \le \alpha \ne \beta \le r} \left(\eta_{\alpha,\beta} \hat{c}^\dagger_\alpha \hat{c}^\dagger_\beta - H.c. \right) \right] \tag{11}$$

over the fermionic raising and lowering operators. Namely

$$\begin{aligned}
\hat{\Omega}^\dagger \hat{c}^\dagger_\alpha \hat{\Omega} = \exp &\left[\sum_{1 \le i \ne j \le r} \left(\eta_{i,j} \hat{c}^\dagger_i \hat{c}^\dagger_j - \eta^*_{i,j} \hat{c}_j \hat{c}_i \right) \right] \hat{c}^\dagger_\alpha \\
&\exp\left[- \sum_{1 \le i \ne j \le r} \left(\eta_{i,j} \hat{c}^\dagger_i \hat{c}^\dagger_j - \eta^*_{i,j} \hat{c}_j \hat{c}_i \right) \right]
\end{aligned} \tag{12}$$

which, exploiting the BCH formula (4) can be written as

$$\hat{\Omega}^\dagger \hat{c}^\dagger_\alpha \hat{\Omega} = \sum_m \frac{1}{m} \left[\sum_{ij} \eta_{i,j} \hat{c}^\dagger_i \hat{c}^\dagger_j - \eta^*_{i,j} \hat{c}_j \hat{c}_i \right]_m \ . \tag{13}$$

It can be shown that in the last summation the two first terms only survive, leading to to the following expression for $\hat{\Omega} \hat{c}^\dagger_\alpha \hat{\Omega}^\dagger$ and $\hat{\Omega} \hat{c}_\gamma \hat{\Omega}^\dagger$ respectively

$$\hat{\Omega}^\dagger \hat{c}^\dagger_\alpha \hat{\Omega} = \hat{c}^\dagger_\alpha + \sum_i \zeta^*_{i\alpha} \hat{c}_i \,,$$

$$\hat{\Omega}^\dagger \hat{c}_\gamma \hat{\Omega} = \hat{c}_\gamma + \sum_i \zeta_{m\gamma} \hat{c}^\dagger_m$$

with $\zeta_{ij} = 2\eta_{ij}$.

We are now in the position to evaluate

$$\mathcal{H}_{cl} = \langle \phi | \hat{\mathsf{H}} - \mu \hat{N} | \phi \rangle \,, \tag{14}$$

where the term $\mu\hat{N}$ has been added to take into account the particle number constraint.

With Eq. (10), Eq. (14) becomes

$$\mathcal{H}_{cl} = \langle 0|\Omega \left[\left(\hat{\mathsf{H}}_0 + \hat{\mathsf{H}}_I\right)\right] \Omega^\dagger|0\rangle \,, \tag{15}$$

where

$$\begin{aligned} \hat{\mathsf{H}}_0 &= \sum_{\alpha,\beta} \Gamma_{\alpha,\beta} \hat{c}^\dagger_\alpha \hat{c}_\beta \,, \\ \hat{H}_I &= \sum_{\alpha,\beta,\gamma,\delta} U_{\alpha,\beta,\gamma,\delta} \hat{c}^\dagger_\alpha \hat{c}^\dagger_\beta \hat{c}_\delta \hat{c}_\gamma \,, \end{aligned}$$

with $\Gamma_{\alpha,\beta} = \lambda_\alpha \delta_{\alpha,\beta} - T_{\alpha,\beta} - \mu\delta_{\alpha,\beta}$. Since Ω is a unitary operator, we can write

$$\begin{aligned} \hat{\Omega}^\dagger \hat{H}_0 \hat{\Omega} &= \sum_{\alpha,\beta} \Gamma_{\alpha,\beta} \hat{\Omega}^\dagger \hat{c}^\dagger_\alpha \hat{\Omega}\hat{\Omega}^\dagger \hat{c}_\beta \hat{\Omega} \,, \\ \hat{\Omega}^\dagger \hat{H}_I \hat{\Omega} &= \sum_{\alpha,\beta,\gamma,\delta} U_{\alpha,\beta,\gamma,\delta} \hat{\Omega}^\dagger \hat{c}^\dagger_\alpha \hat{\Omega}\hat{\Omega}^\dagger \hat{c}^\dagger_\beta \hat{\Omega}\hat{\Omega}^\dagger \hat{c}_\delta \hat{\Omega}\hat{\Omega}^\dagger \hat{c}_\gamma \hat{\Omega} \,. \end{aligned} \tag{16}$$

The following expectation values must then be evaluated. For the one-body term

$$\hat{\Omega}^\dagger \hat{H}_0 \hat{\Omega} = \sum_{\alpha,\beta} \Gamma_{\alpha,\beta} \left[\hat{c}^\dagger_\alpha + \sum_i \zeta^*_{i\alpha} \hat{c}_i\right] \cdot \left[\hat{c}_\beta + \sum_k \zeta_{k\beta} \hat{c}^\dagger_k\right] \tag{17}$$

and for the interaction term

$$\begin{aligned} \hat{\Omega}^\dagger \hat{H}_I \hat{\Omega} = \sum_{\alpha,\beta,\gamma,\delta} U_{\alpha,\beta,\gamma,\delta} &\left[\hat{c}^\dagger_\alpha + \sum_i \zeta^*_{i\alpha} \hat{c}_i\right] \cdot \left[\hat{c}^\dagger_\beta + \sum_j \zeta^*_{j\beta} \hat{c}_j\right] \cdot \\ &\left[\hat{c}_\delta + \sum_k \zeta_{k\delta} \hat{c}^\dagger_k\right] \cdot \left[\hat{c}_\gamma + \sum_l \zeta_{l\gamma} \hat{c}^\dagger_l\right] \,. \end{aligned} \tag{18}$$

As it can be directly verified, in the calculation of the expectation values over the vacuum state $|0\rangle$ only the following terms survive

$$\langle 0|\hat{\Omega}^\dagger \hat{H}_0 \hat{\Omega}|0\rangle = \sum_{\alpha,\beta} \sum_{ij} \Gamma_{\alpha,\beta} \zeta^*_{i\alpha} \zeta_{j\beta} \langle 0|\hat{c}_i \hat{c}^\dagger_j|0\rangle \,, \tag{19}$$

$$\begin{aligned} \langle 0|\hat{\Omega}^\dagger \hat{H}_I \hat{\Omega}|0\rangle = \sum_{\alpha,\beta,\gamma,\delta} U_{\alpha,\beta,\gamma,\delta} &\Bigg(\sum_{i,j,k,l} \zeta^*_{i\alpha} \zeta^*_{j\beta} \zeta_{k\delta} \zeta_{l\gamma} \langle 0|\hat{c}_i \hat{c}_j \hat{c}^\dagger_k \hat{c}^\dagger_l|0\rangle \\ &+ \sum_{i,j} \zeta^*_{i\alpha} \zeta_{j\gamma} \langle 0|\hat{c}_i \hat{c}^\dagger_\beta \hat{c}_\delta \hat{c}^\dagger_j|0\rangle \Bigg) \,. \end{aligned} \tag{20}$$

The two expectation values over the vacuum state give

$$\langle 0|\hat{c}_i \hat{c}^\dagger_j|0\rangle = \delta_{ij} \tag{21}$$

$$\langle 0|\hat{c}_i \hat{c}_j \hat{c}^\dagger_l \hat{c}^\dagger_k|0\rangle = \delta_{il}\delta_{jk} - \delta_{jk}\delta_{il} \tag{22}$$

$$\langle 0|\hat{c}_i \hat{c}^\dagger_\beta \hat{c}_\delta \hat{c}^\dagger_j|0\rangle = \delta_{i\beta}\delta_{j\delta} \tag{23}$$

hence

$$\mathcal{H}^{cl} = \sum_{\alpha\beta} \Gamma_{\alpha\beta} \sum_i \zeta^*_{i\alpha} \zeta_{i\beta} + \sum_{\alpha,\beta,\gamma,\delta} U_{\alpha,\beta,\gamma,\delta} \left[\sum_{i,j} \left(\zeta^*_{i\alpha} \zeta^*_{j\beta} \zeta_{j\delta} \zeta_{i\gamma} - \zeta^*_{i\alpha} \zeta^*_{j\beta} \zeta_{i\delta} \zeta_{j\gamma}\right) + \zeta^*_{\beta\alpha} \zeta_{\delta\gamma}\right] \tag{24}$$

4. Effective Hamiltonian

Hamiltonian (24) can be shown to represent the effective Hamiltonian associated with $\hat{H}$ within the time-dependent variational principle procedure (4). The latter is based on approximating the quantum states of the system by a trial state $|\Psi\rangle$ satisfying the weak form of the Schrödinger equation $\langle\Psi|i\hbar\partial_t - \hat{H}|\Psi\rangle = 0$. Here, we assume that $|\Psi\rangle$, up to an irrelevant phase factor, is the coherent state defined in equation (10). The variational procedure allows one to derive the effective Lagrangian $\dot{S} = \langle\Psi|i\hbar\partial_t - \hat{H}|\Psi\rangle$, depending on dynamical variables $\zeta_{\alpha\beta}$, which in turn supplies the effective Hamiltonian (24). Such a procedure provides as well the dynamical equations pertaining to Hamiltonian (24) and the relevant Lie-Poisson brackets. The latter exhibit the same algebraic structure of commutators (6) and will be defined below.

A quite direct physical insight about coherent-state parameters $\zeta_{\alpha\beta}$ is achieved when considering the expectation values for the elements of the Lie algebra so(2r) over the coherent states $|\phi\rangle$. We have

$$\langle\phi|\hat{c}^\dagger_\alpha\hat{c}^\dagger_\beta|\phi\rangle = \zeta^*_{\beta\alpha}\,, \tag{25}$$

$$\langle\phi|\hat{c}^\dagger_\alpha\hat{c}_\beta|\phi\rangle = \sum_i \zeta^*_{i\alpha}\zeta_{i\beta} = \xi_{\alpha\beta}\,, \tag{26}$$

$$\langle\phi|\hat{c}_\alpha\hat{c}_\beta|\phi\rangle = \zeta_{\alpha\beta}\,, \tag{27}$$

showing how parameters $\zeta_{\alpha\beta}$ are related to microscopic physical processes of creation/destruction of lattice fermions. Moreover Eq. (24) can be written as

$$\mathcal{H}_{cl} = \sum_{\alpha\beta}\Gamma_{\alpha\beta}\xi_{\alpha\beta} + \sum_{\alpha,\beta,\gamma,\delta} U_{\alpha,\beta,\gamma,\delta}\left[(\xi_{\alpha\gamma}\xi_{\beta\delta} - \xi_{\alpha\delta}\xi_{\beta\gamma}) + \zeta^*_{\beta\alpha}\zeta_{\delta\gamma}\right]\,. \tag{28}$$

The three terms represent the direct, the exchange and the pairing term in the HFB mean-field approximation.

4.1. Evolution Equations for the canonical variables

According to (4) the variables $\zeta_{\alpha\beta}$, $\zeta^*_{\alpha\beta}$ and $\xi_{\alpha\beta}$ represent the canonical variable for the classical Hamiltonian $\mathcal{H}_{cl}$. With a well-known procedure (10) it is possible to describe the time evolution of those canonical variables in terms of their Poisson brackets with $\mathcal{H}_{cl}$. To write the Lie-Poisson brackets for the given dynamical system we can make explicit the structure constants for the so(2r) algebra

$$\begin{aligned}
c_{1,\alpha,\beta;1,\gamma,\delta}{}^{q,\mu,\nu} &= c_{2,\alpha,\beta;2,\gamma,\delta}{}^{q,\mu,\nu} = 0\,,\\
c_{1,\alpha,\beta;2,\gamma,\delta}{}^{q,\mu,\nu} &= \delta_{q,3}\left(\delta_{\mu\gamma}\delta_{\nu\beta}\delta_{\alpha\delta} + \delta_{\mu\delta}\delta_{\nu\alpha}\delta_{\beta\gamma} - \delta_{\mu\gamma}\delta_{\nu\alpha}\delta_{\beta\delta} - \delta_{\mu\delta}\delta_{\nu\beta}\delta_{\alpha\gamma}\right)\,,\\
c_{1,\alpha,\beta;3,\gamma,\delta}{}^{q,m,n} &= \delta_{q,1}\left(\delta_{\mu\alpha}\delta_{\nu\delta}\delta_{\beta\gamma} - \delta_{\mu\beta}\delta_{\nu\delta}\delta_{\alpha\gamma}\right)\,,\\
c_{2,\alpha,\beta;3,\gamma,\delta}{}^{q,m,n} &= \delta_{q,2}\left(\delta_{\mu\beta}\delta_{\nu\gamma}\delta_{\alpha\delta} - \delta_{\mu\alpha}\delta_{\nu\gamma}\delta_{\beta\delta}\right)\,.
\end{aligned} \tag{29}$$

Thus the Poisson brackets have the following form

$$
\begin{aligned}
\{f,g\} = \quad & \sum_{\alpha\beta\gamma\delta} \left(\xi_{\gamma\beta}\delta_{\alpha\gamma} - \xi_{\gamma\alpha}\delta_{\beta\delta} + \xi_{\delta\alpha}\delta_{\beta\gamma} - \xi_{\delta\beta}\delta_{\alpha\gamma}\right) \left(\frac{\partial f}{\partial \zeta_{\alpha\beta}} \frac{\partial g}{\partial \zeta^*_{\gamma\delta}} - \frac{\partial f}{\partial \zeta^*_{\gamma\delta}} \frac{\partial g}{\partial \zeta_{\alpha\beta}} \right) + \\
& + \left(\zeta_{\alpha\delta}\delta_{\gamma\beta} - \zeta_{\beta\delta}\delta_{\gamma\alpha}\right) \left(\frac{\partial f}{\partial \zeta_{\alpha\beta}} \frac{\partial g}{\partial \xi_{\gamma\delta}} - \frac{\partial f}{\partial \xi_{\gamma\delta}} \frac{\partial g}{\partial \zeta_{\alpha\beta}} \right) + \\
& + \left(\zeta_{\gamma\beta}\delta_{\alpha\delta} - \zeta_{\alpha\gamma}\delta_{\beta\delta}\right) \left(\frac{\partial f}{\partial \zeta^*_{\alpha\beta}} \frac{\partial g}{\partial \xi_{\gamma\delta}} - \frac{\partial f}{\partial \xi_{\gamma\delta}} \frac{\partial g}{\partial \zeta^*_{\alpha\beta}} \right) + \\
& + \left(\xi_{\beta\delta}\delta_{\gamma\alpha} - \xi_{\alpha\gamma}\delta_{\beta\delta}\right) \left(\frac{\partial f}{\partial \xi_{\alpha\beta}} \frac{\partial g}{\partial \xi_{\gamma\delta}} - \frac{\partial f}{\partial \xi_{\gamma\delta}} \frac{\partial g}{\partial \xi_{\alpha\beta}} \right) . \qquad (30)
\end{aligned}
$$

Remembering that

$$
\dot{\zeta}_{\rho,\theta} = \{\zeta_{\rho,\theta}, H\} \ , \qquad (31)
$$

it is possible to write

$$
\begin{aligned}
\dot{\zeta}_{\rho,\theta} = \; & \sum_{\alpha} \left(\Gamma_{\rho\alpha}\zeta_{\theta\alpha} - \Gamma_{\theta\alpha}\zeta_{\rho\alpha}\right) + \\
& \sum_{\alpha\gamma\eta} \left[U_{\alpha\rho\gamma\eta}\left(\zeta_{\theta\gamma}\xi_{\alpha\eta} - \zeta_{\theta\eta}\xi_{\alpha\gamma}\right) + U_{\alpha\theta\gamma\eta}\left(\zeta_{\rho\eta}\xi_{\alpha\gamma} - \zeta_{\theta\gamma}\xi_{\alpha\eta}\right)\right] \\
& \sum_{\beta\gamma\eta} \left[U_{\rho\beta\gamma\eta}\left(\zeta_{\theta\eta}\xi_{\beta\gamma} - \zeta_{\theta\gamma}\xi_{\beta\eta}\right) + U_{\theta\beta\gamma\eta}\left(\zeta_{\rho\eta}\xi_{\beta\gamma} - \zeta_{\rho\gamma}\xi_{\beta\eta}\right)\right] + \\
& \sum_{\alpha\gamma\eta} \zeta_{\eta\gamma}\left(U_{\alpha\theta\gamma\eta}\xi_{\alpha\rho} - U_{\alpha\rho\gamma\eta}\xi_{\alpha\theta}\right) + \\
& \sum_{\beta\gamma\eta} \zeta_{\eta\gamma}\left(U_{\rho\beta\gamma\eta}\xi_{\beta\theta} - U_{\theta\rho\gamma\eta}\xi_{\beta\rho}\right) \qquad (32)
\end{aligned}
$$

that provide the set of dynamical equations governing the evolution of the coherent state that approximates the system quantum state. In particular, they allow to find the mean-field ground the state for the system and to perform a weakly-excited state analysis.

5. Conclusions

In this paper we have formulated an HFB mean-field approximation for a one-dimensional array of oblate Harmonic Oscillators loaded with neutral fermionic atoms. As already pointed out by Grasso et al. (8), the numerical solution to Eq. (32) appears to be rather demanding from a computational point of view. It seems then appropriate for future work to concentrate on the simplest situation beyond known models like, as already mentioned, a dimer with a six-level local structure.

In this case the evaluation of ground-state properties in this mean-field picture as a function of the relevant parameters(i.e $T_{\alpha,\beta}$, $U_{\alpha\beta\gamma\delta}$, μ) reduces to the fixed-point analysis of Eq.(32). Moreover, an extension to finite-temperature properties does not seem beyond the possibilities of the analytical techniques here outlined and may represent one of the future lines of research.

Appendix A. Expectation values calulation

In the present section we will explicitly calculate the terms obtained form Eqs. (17,18) leading to Eqs. (19). To evaluate (17) we need to perform the following product calculation

$$\hat{\Omega}^\dagger \hat{H}_0 \hat{\Omega} = \sum_{\alpha\beta ij} \Gamma_{\alpha,\beta} \left[\hat{c}^\dagger_\alpha + \sum_i \zeta^*_{i\alpha} \hat{c}_i \right] \cdot \left[\hat{c}_\beta + \sum_j \zeta_{j\beta} \hat{c}^\dagger_j \right] \tag{A.1}$$

which is equal to

$$\hat{\Omega}^\dagger \hat{H}_0 \hat{\Omega} = \sum_{\alpha\beta} \left[\hat{c}^\dagger_\alpha \hat{c}_\beta + \sum_i \zeta^*_{i\alpha} \hat{c}_i \hat{c}_\beta + \sum_j \zeta_{j\beta} \hat{c}^\dagger_\alpha \hat{c}^\dagger_j + \sum_{ij} \zeta^*_{i\alpha} \zeta_{j\beta} \hat{c}_i \hat{c}^\dagger_j \right]. \tag{A.2}$$

The evaluation of Eq. (A.2) over the vacuum state $|0\rangle$ leads to vanishing contributions for all non number-conseving terms and for all the terms with a lowering operator on the right-hand side (or a raising operator on the left-hand side). Namely (see Eq. (19)

$$\langle 0 | \hat{\Omega}^\dagger \hat{H}_0 \hat{\Omega} | 0 \rangle = \sum_{\alpha\beta} \sum_{ij} \Gamma_{\alpha,\beta} \zeta^*_{i\alpha} \zeta_{j\beta} \langle 0 | \hat{c}_i \hat{c}^\dagger_j | 0 \rangle. \tag{A.3}$$

With an analogous procedure it is possible to evaluate the expression given by Eq. (18)

$$\begin{aligned} \hat{\Omega}^\dagger \hat{H}_I \hat{\Omega} = \sum_{\alpha,\beta,\gamma,\delta} U_{\alpha,\beta,\gamma,\delta} \left[\hat{c}^\dagger_\alpha + \sum_i \zeta^*_{i\alpha} \hat{c}_i \right] \cdot \left[\hat{c}^\dagger_\beta + \sum_j \zeta^*_{j\beta} \hat{c}_j \right] \cdot \\ \left[\hat{c}_\delta + \sum_k \zeta_{k\delta} \hat{c}^\dagger_k \right] \cdot \left[\hat{c}_\gamma + \sum_l \zeta_{l\gamma} \hat{c}^\dagger_l \right] \end{aligned} \tag{A.4}$$

leading to

$$\begin{aligned} \hat{\Omega}^\dagger \hat{H}_I \hat{\Omega} = \sum_{\alpha,\beta,\gamma,\delta} U_{\alpha,\beta,\gamma,\delta} \\ \Bigg[\hat{c}^\dagger_\alpha \hat{c}^\dagger_\beta \hat{c}_\delta \hat{c}_\gamma + \sum_k \zeta_{k\delta} \hat{c}^\dagger_\alpha \hat{c}^\dagger_\beta \hat{c}^\dagger_k \hat{c}_\gamma + \sum_l \zeta_{l\gamma} \hat{c}^\dagger_\alpha \hat{c}^\dagger_\beta \hat{c}_\delta \hat{c}^\dagger_l + \sum_{kl} \zeta_{l\gamma} \zeta_{k\delta} \hat{c}^\dagger_\alpha \hat{c}^\dagger_\beta \hat{c}^\dagger_k \hat{c}^\dagger_l + \\ \sum_j \zeta^*_{j\beta} \hat{c}^\dagger_\alpha \hat{c}_j \hat{c}_\delta \hat{c}_\gamma + \sum_{jk} \zeta^*_{j\beta} \zeta_{k\delta} \hat{c}^\dagger_\alpha \hat{c}_j \hat{c}^\dagger_k \hat{c}_\gamma + \sum_{jl} \zeta^*_{j\beta} \zeta_{l\gamma} \hat{c}^\dagger_\alpha \hat{c}_j \hat{c}_\delta \hat{c}^\dagger_l + \sum_{jkl} \zeta^*_{j\beta} \zeta_{l\gamma} \zeta_{k\delta} \hat{c}^\dagger_\alpha \hat{c}_j \hat{c}^\dagger_k \hat{c}^\dagger_l + \\ \sum_i \zeta^*_{i\alpha} \hat{c}_i \hat{c}^\dagger_\beta \hat{c}_\delta \hat{c}_\gamma + \sum_{ik} \zeta^*_{i\alpha} \zeta_{k\delta} \hat{c}_i \hat{c}^\dagger_\beta \hat{c}^\dagger_k \hat{c}_\gamma + \sum_{il} \zeta^*_{i\alpha} \zeta_{l\gamma} \hat{c}_i \hat{c}^\dagger_\beta \hat{c}_\delta \hat{c}^\dagger_l + \sum_{ikl} \zeta^*_{i\alpha} \zeta_{l\gamma} \zeta_{k\delta} \hat{c}_i \hat{c}^\dagger_\beta \hat{c}^\dagger_k \hat{c}^\dagger_l + \\ \sum_{ij} \zeta^*_{i\alpha} \zeta^*_{j\beta} \hat{c}_i \hat{c}_j \hat{c}_\delta \hat{c}_\gamma + \sum_{ijk} \zeta^*_{i\alpha} \zeta^*_{j\beta} \zeta_{k\delta} \hat{c}_i \hat{c}_j \hat{c}_\gamma \hat{c}^\dagger_k + \sum_{ijl} \zeta^*_{i\alpha} \zeta^*_{j\beta} \zeta_{l\gamma} \hat{c}_i \hat{c}_j \hat{c}_\delta \hat{c}^\dagger_l + \\ \sum_{ijkl} \zeta^*_{i\alpha} \zeta^*_{j\beta} \zeta_{k\delta} \zeta_{l\gamma} \hat{c}_i \hat{c}_j \hat{c}^\dagger_k \hat{c}^\dagger_l \Bigg]. \end{aligned} \tag{A.5}$$

With the same argument needed to obtain Eq. (A.3) we can write the expectation value of the operator defined by Eq. (A.5) over the vacuum state $|0\rangle$ as

$$\langle 0 | \hat{\Omega}^\dagger \hat{H}_I \hat{\Omega} | 0 \rangle = \sum_{\alpha,\beta,\gamma,\delta} U_{\alpha,\beta,\gamma,\delta} \Bigg(\sum_{i,j,k,l} \zeta^*_{i\alpha} \zeta^*_{j\beta} \zeta_{k\delta} \zeta_{l\gamma} \langle 0 | \hat{c}_i \hat{c}_j \hat{c}^\dagger_k \hat{c}^\dagger_l | 0 \rangle +$$

$$+ \sum_{i,j} \zeta_{i\alpha}^{*} \zeta_{j\gamma} \langle 0 | \hat{c}_i \hat{c}_\beta^\dagger \hat{c}_\delta \hat{c}_j^\dagger | 0 \rangle \Bigg) \tag{A.6}$$

which is the expression given by Eq. (20).

References

[1] F. Massel and V. Penna. Hubbard-like Hamiltonian for ultracold atoms in a one-dimensional optical lattice. *Phys. Rev. A*, 72:053619, 2005.

[2] S. Peil, J. V. Porto, B. Laburthe Tolra, J. M. Obrecht, B. E. King, 2 S. L. Rolston M. Subbotin, and W. D. Phillips. Patterned loading of a bose-einstein condensate into an optical lattice. *Phys. Rev. A*, 67:051603(R), 2003.

[3] S. J. J. M. F. Kokkelmans, J. N. Milstein, M. L. Chiofalo, R. Walser, and M. J. Holland. Resonance superfluidity: Renormalization of resonance scattering theory. *Phys. Rev. A*, 65:053617, 2002.

[4] WM Zhang, DH Feng, and R. Gilmore. Coherent states: Theory and some applications. *Rev. Mod. Phys*, 62:867–927, 1990.

[5] Montorsi A., Rasetti M., and Solomon A. I. Dynamical Superalgebra and Supersymmetry for a Many-Fermion System. *Phys. Rev. Lett.*, 59(20):2244, 1987.

[6] A. Montorsi and V. Penna. Spin picture of the one-dimensional Hubbard model: Two-fluid structure and phase dynamics. *Phys. Rev. B*, 60:12069, 1999.

[7] V. Bach, EH Lieb, and JP Solovej. Generalized Hartree-Fock Theory and the Hubbard Model. *J. Stat. Phys.*, 76:3–90, 1994.

[8] Grasso M. and Urban M. Hartree-Fock-Bogoliubov theory versus local-density approximation for superfluid trapped fermionic atoms. *Phys. Rev. A*, 68:033610, 2003.

[9] Alexander Albus, Fabrizio Illuminati, and Jens Eisert. Mixtures of bosonic and fermionic atoms in optical lattices. *Phys. Rev. A*, 68:023606, 2003.

[10] A. Perelomov. *Generalized Coherent States and Their Applications.* Springer, Berlin, 1986.